

<b>Antibiótico</b>	<b>Dosis</b>	<b>Espectro</b>
Cefalexina	20-30 mg/kg/8 h	<i>S.intermedius</i>
Cefazolina	20 mg/kg/6 h	“
Cloxacilina	10 mg/kg/6 h	“
Amoxicilina	20 mg/kg/12 h	<i>Streptococos sps.</i>
Enrofloxacina	10-20 mg/kg/24 h	<i>Brucella canis</i>
Doxiciclina	25 mg/kg/24 h	“
Gentamicina	5 mg/kg/24 h	“
Penicilina G	40 mg/kg/12 h	<i>Actinomyces sps.</i>
Penicilina G	100.000 U/kg/ 6h	“
Ketoconazole	10 mg/kg/12 h	<i>Coccidioides immitis</i>
Fluconazole	5 mg/kg/12 h	“
Amfotericina B	0.25 mg/kg/ 48 h	<i>Aspergillus sps.</i>
Itraconazole	5 mg/kg/ 12 o 24 h	“



Por: Xavier Raurell.

### **Tabla de antibióticos usados en discospondilitis.**

Tabla extraída de W.Thomas.Diskospondylitis and Other Vertebral Infections.The Veterinary Clinics of North America,Vol.30, nº1, Enero, 2000.